

# THE MOMENTS OF THE NUMBER OF CYCLES OF A RANDOM PERMUTATION BY SIMPLE ENUMERATION

By T. LENGYEL  
*Occidental College, Los Angeles*

*SUMMARY.* We present a new proof for the Poisson limit distribution of the number of fixed points of a random permutation. Despite the combinatorial nature of the proof, it does not involve the use of inclusion-exclusion principle, cycle representation of permutations, number of derangements, rewriting formulas for the distribution of the fix points, generating functions, or transformation formulas between moments. The proof is elementary in terms of enumeration and based on the notion of Stirling numbers. It requires some familiarity with the moment generating function of the Poisson distribution and the Fréchet-Shohat moment convergence theorem. The method is extended to the distribution of the number of  $k$ -cycles of an  $n$ -element set, for  $k \leq n$ .

## 1. Introduction

Let  $X_{n,r}$  denote the number of  $r$ -cycles of a random permutation over an  $n$ -element set  $[n] = \{1, 2, \dots, n\}$ . We use the convenient  $X_n = X_{n,1}$  notation for the number of fixed points. Let  $p_n(k)$  denote the probability that a random permutation over  $[n]$ , for short a random  $n$ -permutation, has  $k$  fixed points, i.e.,  $p_n(k) = P(X_n = k)$ . We find the  $k$ -th moments  $M_k(n) = E(X_n^k)$  and  $M_k(n, r) = E(X_{n,r}^k)$  without using any inclusion-exclusion, inversion or moment transformation formula or determining the related probabilities.

The convergence of the distribution of  $X_n$  to the Poisson distribution is well known. The limiting distribution can be determined by using formulas for  $p_n(k)$ . We take a different approach and find the limit distributions (Corollary) based on a moment convergence theorem by calculating the moments (Theorems 1 and 2). Although we might encounter difficulties for other distributions, in this case we can give a simple interpretation of the moments without determining the explicit probabilities.

---

Paper received. May 1993. revised June 1996.

AMS (1991) subject classifications. 60C05, 05A05, 60E10, 05A19.

Key words and phrases. Random permutations, moments, combinational probability.

We note that Takács (1991) has recently proved a general convergence theorem based on the binomial moments which in addition yields an *explicit* formula for the limit distribution. We will use neither binomial nor factorial moments.

In this paper  $S(n, k)$  denotes the Stirling number of the second kind, i.e., the number of partitions of  $n$  distinct elements into  $k$  non-empty subsets and  $(x)_k$  stands for  $x(x-1)\dots(x-k+1)$  where  $k = 1, 2, \dots$ . The Bell number,  $\omega(n)$ , is defined as the number of all partitions of  $[n]$ , i.e.,  $\omega(n) = \sum_{k=1}^n S(n, k)$ .

## 2. The number of fixed points and $k$ -cycles of a random permutation

The problem of analyzing  $p_n(k)$  and the expected value of the number of fixed points of a random  $n$ -permutation is probably originated in card games. We count the matches when the cards of two well-shuffled decks are matched against each other. The first solution goes back to Montmort who found a recurrence relation for the number of matches in 1708. The problem has been generalized in many ways and counting unrestricted and restricted permutations of an  $n$ -element set has been a popular area of research (e.g., Penrice (1991) and Takács (1981)).

The probability  $p_n(k)$  is usually determined by the inclusion-exclusion principle (cf. Feller (1968), Graham et al. (1988), and Wilf (1990)). Notice that  $p_n(n-1) = 0$ . It also follows that  $p_n(0) = \sum_{i=0}^n (-1)^i \frac{1}{i!} = e^{-1} + O(1/n!)$  and  $p_n(k) = \frac{1}{k!} \sum_{i=0}^{n-k} (-1)^i \frac{1}{i!}$ , which yields the asymptotic formula  $p_n(k) = \frac{1}{ek!} + O(\frac{1}{(n-k)!})$ , for every fixed  $k$  (see e.g., Feller (1968)). The asymptotic behavior of  $p_n(k)$  can be derived by the method of generating functions (cf. Wilf (1990)) too. Roughly speaking,  $p_n(k)$  conforms to the Poisson law with parameter 1 as  $n$  tends to  $\infty$ . Similar results hold for the distribution of  $X_{n,r}$ . We give an alternative proof of these facts based on the moments.

Observe that the probabilities are nearly independent of  $n$ . It turns out that for  $n \geq 2$ ,  $M_1(n) = E(X_n) = 1$  and  $M_2(n) = E(X_n^2) = 2$  do not depend on  $n$ . In general, one can calculate  $M_k(n)$  by

**THEOREM 1.** *The moments  $M_k(n) = E(X_n^k)$  of the number of fixed points of a random  $n$ -permutation are*

$$M_k(n) = \begin{cases} \omega(k), & \text{if } k \leq n \\ \sum_{i=1}^n S(k, i) & \text{if } k > n. \end{cases}$$

Note that for every  $n$ , the first  $n$  moments of  $X_n$  do not depend on  $n$ . We set  $M_0(n) = 1 (n \geq 1)$ .

**PROOF.** Let  $Y_i$  be the indicator random variable of the event that element  $i$  is a fixed point of the random  $n$ -permutation, i.e.,

$$Y_i = \begin{cases} 1, & \text{if element } i \text{ is a fixed point,} \\ 0, & \text{otherwise.} \end{cases}$$

Clearly, the random variables  $Y_i$  are identically distributed though they are not independent. For the number of fixed points  $X_n$ , we get  $X_n = \sum_{i=1}^n Y_i$ . We evaluate  $M_k(n) = E(X_n^k)$  by expanding the right hand side of this equation using the multinomial expansion. That is,

$$M_k(n) = E((Y_1 + Y_2 + \dots + Y_n)^k) = \sum_{\mathcal{N}_n} \binom{k}{i_1 \dots i_n} E(Y_1^{i_1} Y_2^{i_2} \dots Y_n^{i_n})$$

where the summation extends over the set  $\mathcal{N}_n$  of  $n$ -tuples  $(i_1, i_2, \dots, i_n)$  with non-negative integer coordinates provided  $\sum_{j=1}^n i_j = k$ .

Let  $l = l(i_1, i_2, \dots, i_n)$  denote the number of positive exponents in  $Y_1^{i_1} Y_2^{i_2} \dots Y_n^{i_n}$  i.e.,  $l = |\{j | i_j \geq 1\}|$ . Clearly,  $1 \leq l \leq \min\{k, n\}$ . Notice that  $Y_i^c = Y_i$  if  $c \neq 0$ . One can easily see that  $Y_1^{i_1} Y_2^{i_2} \dots Y_n^{i_n}$  and  $Y_1 Y_2 \dots Y_l$  are identically distributed. There are  $\binom{n}{l}$  ways to choose the  $l$  variables with a given set of positive exponents. For  $Y_1 Y_2 \dots Y_l = 1$  if and only if the  $n$ -permutation has the first  $l$  elements of  $[n]$  as fixed points, thus  $E(Y_1 Y_2 \dots Y_l) = \frac{(n-l)!}{n!}$ . Let  $\mathcal{N}_l^+$  denote the set of  $l$ -tuples  $(i_1, i_2, \dots, i_l)$  such that  $i_j \geq 1$ , for all  $j, 1 \leq j \leq l$ , and  $\sum_{j=1}^l i_j = k$ . It follows that

$$M_k(n) = \sum_{l=1}^{\min\{k,n\}} \sum_{\mathcal{N}_l^+} \binom{k}{i_1 \dots i_l} \binom{n}{l} \frac{(n-l)!}{n!} = \sum_{l=1}^{\min\{k,n\}} \sum_{\mathcal{N}_l^+} \binom{k}{i_1 \dots i_l} \frac{1}{l!}.$$

Now observe that

$$\sum_{\mathcal{N}_l^+} \binom{k}{i_1 \dots i_l} = S(k, l) l!$$

for the number of partitions of a  $k$ -element set into  $l$  non-empty blocks is counted on both sides. The order of the blocks is taken into account but not the order of the elements inside the blocks. We find that  $M_k(n) = \sum_{l=1}^{\min\{k,n\}} S(k, l)$ . The proof is now complete. □

REMARK. The previous statement can be derived by using factorial moments for expressing  $M_k(n) = E(X_n^k), k \geq 1$ . In fact, the transformation formula between central and factorial moments,  $X^k = \sum_{l=1}^k S(k, l)(X)_l$ , becomes  $E(X^k) = \sum_{l=1}^k S(k, l) E((X)_l) = \sum_{l=1}^{\min\{k,n\}} S(k, l)$  after taking the expected values. The proof of the formula and the calculation of  $E((X)_l)$ , however, are usually based on the combinatorial theory of the Stirling numbers of the second kind and the number of derangements. Our proof completely avoided this approach. Nonetheless, the identity  $x^k = \sum_{l=1}^k S(k, l)(x)_l, k \geq 1$ , follows easily when the moment enumeration method is applied to binomially distributed random variables.

We can generalize the previous theorem and obtain

**THEOREM 2.** *The moments  $M_k(n, r) = E(X_{n,r}^k)$  of the number of  $r$ -cycles of a random  $n$ -permutation are*

$$M_k(n, r) = \begin{cases} \sum_{i=1}^k S(k, i)/r^i, & \text{if } k \leq [n/r] \\ \sum_{i=1}^{[n/r]} S(k, i)/r^i, & \text{if } k > [n/r]. \end{cases}$$

We note that the  $k$ -th moment of the Poisson distribution with parameter  $\lambda = 1/r$  is  $\sum_{i=1}^k S(k, i)/r^i$ . Theorem 2 and the Fréchet-Shohat moment convergence theorem (see e.g., Moran (1968)) implies that

**COROLLARY.** *The number of  $r$ -cycles of a random  $n$ -permutation has Poisson limit distributions with parameter  $1/r$ .*

**PROOF OF THEOREM 2.** The proof is based on the revised definition of the indicator variables  $Y_i$ . We set  $N = \binom{n}{r}$  and define  $Y_i, i = 1, 2, \dots, N$ , as the indicator variable of the event that the  $i$ -th element in the list of  $r$ -element subsets of  $[n]$  forms an  $r$ -cycle of the random  $n$ -permutation. That is,

$$Y_i = \begin{cases} 1, & \text{if the } i\text{-th } r\text{-element subset of } [n] \text{ is an } r\text{-cycle,} \\ 0, & \text{otherwise.} \end{cases}$$

Let  $l = l(i_1, i_2, \dots, i_n)$  denote the number of positive exponents in  $Y_1^{i_1}, Y_2^{i_2}, \dots, Y_N^{i_N}$ . Incidentally,  $Y_1^{i_1}, Y_2^{i_2}, \dots, Y_N^{i_N}$  is zero or one, and it is one if and only if the corresponding  $l$   $r$ -element subsets form disjoint  $r$ -cycles of the permutation. Provided the subsets are disjoint, there  $((r-1)!)^l (n-lr)!$  ways of forming the required  $r$ -cycles and completing the  $n$ -permutation. We can select the  $l$  disjoint subsets in  $\binom{n}{r \dots r} / l!$  ways (with  $l$  copies of  $r$  under  $n$ ), and there are  $S(k, l)l!$  ways of making  $l$ -tuples with positive coordinates of sum  $k$ . In other words, the multinomial expansion of  $E((Y_1 + Y_2 + \dots + Y_N)^k)$  has  $S(k, l)l!$  terms  $Y_1^{i_1} Y_2^{i_2} \dots Y_N^{i_N}$ , with  $l$  positive exponents, each contributing

$$\binom{n}{r \dots r} \frac{1}{l!} \frac{((r-1)!)^l (n-lr)!}{n!} = \frac{1}{r^l l!}$$

to  $M_k(n, r)$ . Summation on  $l, 1 \leq l \leq \min\{k, [n/r]\}$ , yields the identity for  $M_k(n, r)$ . □

## References

- FELLER, W (1968). *An Introduction to Probability Theory and Its Applications*, vol. I, Wiley, 3rd edition.
- GRAHAM, R.L., KNUTH, D.E. and PATASHNIK, O. (1988). *Concrete Mathematics*, Addison-Wesley, Reading, MA.
- MORAN, P.A.P. (1968). *An Introduction to Probability Theory*, Oxford.
- PENRICE, S.G. (1991). Derangements, permanents, and Christmas presents, *Amer. Math. Monthly*, **94**, 617-620.
- TAKÁCS, L. (1981). On the "Problème des mééges", *Discrete Mathematics*, **36**, 289-297.
- — — (1991). A moment convergence theorem, *Amer. Math. Monthly*, **94**, 742-746.
- WILF, H.S. (1990). *generatortfunctionology*, Academic Press, Boston.

DEPARTMENT OF MATHEMATICS  
OCCIDENTAL COLLEGE  
1600 CAMPUS ROAD  
LOS ANGELES  
CA 90041  
U.S.A.